

# Three new species of *Hydnophlebia* (Polyporales, Basidiomycota) from the Macaronesian Islands

M. Teresa Telleria<sup>1</sup>, Margarita Dueñas<sup>1</sup>, María P. Martín<sup>1</sup>

<sup>1</sup> Department of Mycology, Real Jardín Botánico, RJB/CSIC, Plaza de Murillo 2. 28014 Madrid, Spain

Corresponding author: M. Teresa Telleria ([telleria@rjb.csic.es](mailto:telleria@rjb.csic.es))

---

Academic editor: K. Hosaka | Received 4 July 2017 | Accepted 18 October 2017 | Published 1 November 2017

---

**Citation:** Telleria MT, Dueñas M, Martín MP (2017) Three new species of *Hydnophlebia* (Polyporales, Basidiomycota) from the Macaronesian Islands. MycoKeys 27: 39–64. <https://doi.org/10.3897/mycokeys.27.14866>

---

## Abstract

The genus *Hydnophlebia* includes two species of wood-inhabiting fungi, *Hydnophlebia chrysorhizon* and *Hydnophlebia omnivora*. Both are characterized by cream to reddish-orange, resupinate basidiome, with hydroid hymenophore, margin with strands, a monomitic hyphal system, tubular to ventricose cystidia and elliptical spores. In this paper, a taxonomic study of *Hydnophlebia*, using morphology and molecular analyses of large subunit nuclear ribosomal DNA (LSU) and the internal transcribed spacer nrDNA operon (ITS), is reported. Three new species, *Hydnophlebia canariensis*, *H. gorgonea* and *H. meloi*, from the Macaronesia bioregion (Canary Islands and Cape Verde Archipelago), are described.

## Keywords

Agaricomycetes, corticioid fungi, phylogeny, taxonomy, Canary Islands, Cape Verde Archipelago

## Introduction

*Hydnophlebia* was erected by Parmasto (1967) to accommodate *Hydnum chrysorhizon* Torr. A few years later, the type species was transferred to *Phanerochaete* P. Karst. (Budington and Gilbertson 1973), and *Hydnophlebia* was neglected for a long time. Ryvarden et al. (2005) reintroduced it with a brief description taken from the original Latin diagnosis: basidioma resupinate, membranous, reddish-orange, hymenophore hydroid with aculei, margin with rhizomorphs, hyphal system monomitic, tubular to



ventricose usually few cystidia, and spores ellipsoid, smooth, and thin-walled. According to Hjortstam and Ryvarden (2009), two species should be included in this genus of wood-inhabiting corticioid fungi: *Hydnophlebia chrysorhizon* (Torr.) Parmasto and *Hydnophlebia omnivora* (Shear) Hjortstam & Ryvarden.

Parmasto (1968) included *Hydnophlebia* in tribe Byssomerulieae (Corticaceae) together with other genera, such as *Byssomerulius* Parmasto, *Chaetoderma* Parmasto, *Crustoderma* Parmasto, and *Phanerochaete*, while Larsson (2007), in his phylogenetic classification for corticioid fungi, included it in Meruliaceae, Polyporales. More recently, Floudas and Hibbett (2015) presented a four gene phylogenetic analysis of phanerochaetoid taxa and confirmed *Phanerochaete* as polyphyletic and *Hydnophlebia* as a genus of its own.

During our survey of corticioid fungi from Macaronesia (Canary Islands and Cape Verde Archipelago), nine hydroid specimens were initially identified as belonging to the genus *Phanerochaete*. BLAST search of the large subunit of the nrDNA (LSU) sequences showed high similarity with a sequence published in Wu et al. (2010) and identified as *Phanerochaete chrysorhizon* (Torr.) Budington & Gilb. (AF139967). In the analysis by Wu et al. (2010) this sequence was recovered within a clade (clade V) containing i.a. *Phlebia* sensu stricto and a number of taxa with typically odontoid or hydroid hymenophore, quite far from the *Phanerochaete* core group. BLAST search of the internal transcribed spacers of the nuclear ribosomal gene (ITS) sequences, which gave high similarity to sequences labelled as *Phanerochaete chrysorhizon* (AY219359) and *Phanerochaete omnivora* (Shear) Burdsall & Nakasone (AY219360) published in de Koker et al. (2003). Like later Wu et al. (2010) also de Koker et al. (2003) found that these taxa were not related to the *Phanerochaete* core group.

The aim of this study was to characterize and classify our specimens from Macaronesia, using morphological data and molecular analyses of LSU and ITS regions.

## Materials and methods

### Sampling, morphological studies and line drawings

Specimens were collected in the Canary Islands and Cape Verde Archipelago (Table 1), and are deposited in the mycological collection (MA-Fungi) of the Real Jardín Botánico herbarium in Madrid, Spain; the initials MD correspond to M. Dueñas, and Tell. to M.T. Telleria. The type specimens of *Hydnum chrysorhizon* (NY!) and *Hydnum omnivorum* Shear (BPI!) were included in the morphological analyses. Colours of dried basidiomata are given according to ISCC-NBS Centroid Color Charts (Kelly and Judd 1976). Dried specimens were also used for light microscope studies and drawings. Measurements and drawings were made from microscopic sections mounted in 3% aqueous KOH and/or Congo Red solution and examined at magnifications up to 1250× using an Olympus BX51 microscope. The length and width of 30 spores and 10 basidia were measured from each sample. Line drawings were made with a Leica DM2500 microscope with the aid of a drawing tube.



## DNA isolation and sequencing

Genomic DNA was extracted from eight collections (Table 1) using DNeasy® Plant Mini Kit (QIAGEN, Valencia, CA), following the manufacturer's instructions. Basidiomes were disrupted using Tissue-Lyser II (QIAGEN, Germany) and glass beads. Lysis buffer incubation was overnight at 55 °C.

Total DNA was used for PCR amplification of the D1–D2 region of the large subunit (LSU) and the internal transcribed spacer region (ITS) of the nuclear ribosomal gene. The primers LR0R (Rehner and Samuels 1994) and LR7 (Vilgalys and Hester 1990) were used to amplify the region of the LSU nrDNA; the primers ITS1F (Gardes and Bruns 1993) and ITS4 (White et al. 1990) were used to obtain amplifications of both ITS regions, including the 5.8S of the ribosomal RNA gene cluster and flanking parts of the small subunit (SSU) and large subunit (LSU) nuclear ribosomal genes. Individual reactions to a final volume of 25 µL were carried out using illustra™ PuReTaq™ Ready-To-Go™ PCR Beads (GE Healthcare, Buckingham) with a 10 pmol/µL primer concentration, following the thermal cycling conditions used in Martín and Winka (2000).

Negative controls lacking fungal DNA were run for each experiment to check for contamination. The reactions were run with the following parameters for the LSU nrDNA: initial denaturation at 94 °C for 5 min, then 36 cycles of denaturation at 94 °C for 30 s, annealing at 52 °C for 30 s, and extension at 72 °C for 1.5 min, with a final extension at 72 °C for 10 min, and 4 °C soak; for the ITS nrDNA: initial denaturation at 95 °C for 5 min, then 5 cycles of denaturation at 95 °C for 30 s, annealing at 54 °C for 30 s, and extension at 72 °C for 1 min, followed by 33 cycles of denaturation at 72 °C for 1 min, annealing at 48 °C for 30 s, and extension at 72 °C, with a final extension at 72 °C for 10 min and 4 °C soak.

PCR products were checked on 2% agarose D1 low EEO (CONDA, Pronadisa™) gels and subsequently purified using the QIAquick Gel PCR Purification (QIAGEN) kit according to the manufacturer's instructions. The purified PCR products were sequenced using the same amplification primers at Macrogen Korea (Seoul, Korea).

Sequencher v. 4.2 (Gene Codes Corporation, Ann Arbor, MI) was used to edit the resulting electropherograms and to assemble contiguous sequences (Table 1 in bold). BLAST searches (Altschul et al. 1997), using the MEGABLAST option were done to compare the sequences obtained against the sequences in the EMBL/GenBank/DDBJ databases (Cochrane et al. 2011, 2016).

## Sequence alignment and phylogenetic analyses

The LSU and ITS sequences obtained were aligned separately using Se-Al v. 2.0a11 Carbon (Rambaut 2002) for multiple sequences.

To infer phylogenetic relationships of Macaronesian specimens within Meruliaceae, the LSU sequences were compared with homologous sequences retrieved from the



**Table 1.** Specimens of *Hydnophlebia* species described as new, and EMBL/GenBank/DDBJ and UNITE accessions included in the LSU and ITS nrDNA analyses. The asterisk (\*) after the taxon names denotes type species of the genus. The specimens with uncertain generic placement are listed at the end of the table; in Fig. 1 and 2, the uncertainty is indicated by brackets around the name. Isolates and/or voucher specimens are indicated as they appear in GenBank and UNITE accessions.

Names after our LSU or ITS analyses	Names included in EMBL/GenBank/DDBJ and UNITE	Isolate/Voucher	GenBank/UNITE accessions	
			LSU	ITS
<i>Abortiporus biennis</i> *	<i>Abortiporus biennis</i>	KEW210	AF287842	–
<i>Cabaladontia queletii</i> *	<i>Phlebia queletii</i>	FCUG 722, culture	AF141626	–
<i>Ceriporia viridans</i> *	<i>Ceriporia viridans</i>	FO24398	AJ406518	–
<i>Ceriporiopsis gilvescens</i> *	<i>Ceriporiopsis gilvescens</i>	O/Haussknecht98	DQ144618	–
<i>Climacodon septentrionalis</i> *	<i>Climacodon septentrionalis</i>	HHB-13438-sp	AF518610	–
<i>Crustodontia chrysocreas</i> *	<i>Phlebia chrysocreas</i>	FPL-6080	AY293199	–
<i>Crustodontia chrysocreas</i> *	<i>Phlebia chrysocreas</i>	KHL10216 (GB)	AY586695	–
<i>Cymatoderma elegans</i> *	<i>Cymatoderma elegans</i>	Halling9064 (NY)	JN649341	–
<i>Hydnophlebia canariensis</i>	<i>Hydnophlebia canariensis</i>	17035Tell., MA-Fungi 86622, Holotype	<b>KF528103</b>	<b>KF483012</b>
<i>Hydnophlebia canariensis</i>	<i>Hydnophlebia canariensis</i>	17038Tell., MA-Fungi 86623	<b>KF528104</b>	<b>KF483013</b>
<i>Hydnophlebia canariensis</i>	<i>Hydnophlebia canariensis</i>	17674Tell., MA-Fungi 86619	<b>KF528100</b>	<b>KF483009</b>
<i>Hydnophlebia chrysorhizon</i> *	<i>Phanerochaete chrysorhizon</i>	FP-102002-sp (CFMR)	–	AY219359
<i>Hydnophlebia chrysorhizon</i> *	<i>Phanerochaete chrysorhizon</i>	T-484, RGT 871020/12	AF139967	–
<i>Hydnophlebia chrysorhizon</i> *	<i>Hydnophlebia chrysorhizon</i>	T 484	–	KP135335
<i>Hydnophlebia chrysorhizon</i> *	<i>Hydnophlebia chrysorhizon</i>	FP-134985	–	KP135336
<i>Hydnophlebia chrysorhizon</i> *	<i>Hydnophlebia chrysorhizon</i>	HHB-18767	–	KP135337
<i>Hydnophlebia chrysorhizon</i> *	<i>Hydnophlebia chrysorhizon</i>	FD-282	–	KP135338
<i>Hydnophlebia gorgonea</i>	<i>Hydnophlebia gorgonea</i>	13327MD, MA-Fungi 86642	<b>KF528122</b>	<b>KF483031</b>
<i>Hydnophlebia gorgonea</i>	<i>Hydnophlebia gorgonea</i>	19110Tell., MA-Fungi 86658	<b>KF528139</b>	<b>KF483048</b>
<i>Hydnophlebia gorgonea</i>	<i>Hydnophlebia gorgonea</i>	19111Tell., MA-Fungi 86659, Holotype	<b>KF528140</b>	<b>KF483049</b>
<i>Hydnophlebia gorgonea</i>	<i>Hydnophlebia gorgonea</i>	19133Tell., MA-Fungi 86664	<b>KF528145</b>	<b>KF483054</b>
<i>Hydnophlebia meloi</i>	<i>Hydnophlebia meloi</i>	19071Tell., MA-Fungi 86654, Holotype	<b>KF528135</b>	<b>KF483044</b>
<i>Hydnophlebia omnivora</i>	<i>Phanerochaete omnivora</i>	HHB-5969-sp	–	AY219360
<i>Hydnophlebia omnivora</i>	<i>Hydnophlebia omnivora</i> 2 <sup>a</sup>	ME-497	–	KP135332
<i>Hydnophlebia omnivora</i>	<i>Hydnophlebia omnivora</i> 2 <sup>a</sup>	HHB-6228-sp	–	KP135333
<i>Hydnophlebia</i> sp. 1	<i>Hydnophlebia omnivora</i> 1 <sup>a</sup>	KKN-112-sp	–	KP135334
<i>Hydnophlebia</i> sp. 2	<i>Phlebia</i> sp.	TU108437	–	UDB016816
<i>Junghuhnia crustacea</i> *	<i>Junghuhnia crustacea</i>	X1127, O. Miettinen 13852,1 (H)	JN710554	–
<i>Junghuhnia crustacea</i> *	<i>Junghuhnia crustacea</i>	X262, O. Miettinen 2954,1 (H)	JN710553	–
<i>Lamelloporus americanus</i> *	<i>Lamelloporus americanus</i>	X670, T. Laessoe 10119 (O, H)	JN710567	–
<i>Lilaceophlebia livida</i> *	<i>Phlebia livida</i>	FCUG 2189, culture	AF141624	–
<i>Merulius tremellosus</i> *	<i>Phlebia tremellosa</i>	FPL-4294	AY293200	–
<i>Merulius tremellosus</i> *	<i>Phlebia tremellosa</i>	FCUG 1813, culture	AF141632	–
<i>Merulius tremellosus</i> *	<i>Phlebia tremellosa</i>	F15198 (UBC)	DQ384584 <sup>b,c</sup>	DQ384584 <sup>b,c</sup>



Names after our LSU or ITS analyses	Names included in EMBL/GenBank/DDBJ and UNITE	Isolate/Voucher	GenBank/UNITE accessions	
			LSU	ITS
<i>Merulius tremellosus</i> *	<i>Phlebia tremellosa</i>	CIRM-BRFM 968	–	GU731568
<i>Mycoacia fuscoatra</i> *	<i>Mycoacia fuscoatra</i>	KHL13275 (GB)	JN649352	JN649352
<i>Mycoacia fuscoatra</i> *	<i>Phlebia subserialis</i>	KUC8041, culture	AY858370	
<i>Mycoacia nothofagi</i>	<i>Mycoacia nothofagi</i>	KHL13750	GU480000	–
<i>Mycoacia nothofagi</i>	<i>Phlebia nothofagi</i>	AH31887	GQ259416	–
<i>Mycoacia nothofagi</i>	<i>Phlebia nothofagi</i>	KHL13750	GU226430	–
<i>Mycorrhaphium adustum</i> *	<i>Mycorrhaphium adustum</i>	KHL12255 (GB)	JN710573	–
<i>Mycoaciella bispora</i> *	<i>Mycoaciella bispora</i>	EL13_99	AY586692	–
<i>Phlebia acerina</i>	<i>Phlebia acerina</i>	FCUG 568, culture	AF141615	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	culture?	AB325676	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	FCUG2423, culture	AF141627	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	FPL6140	AF287885	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	GEL5258	AJ406541	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	KUC8034, culture	AY858369	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	TM03_491	EU522844	–
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	JLL-15608-sp. (CFMR)	–	AY219366
<i>Phlebia radiata</i> *	<i>Phlebia radiata</i>	ATCC 64658, culture	–	EF491867
<i>Phlebia rufa</i>	<i>Phlebia rufa</i>	FCUG 2397	AF141628	–
<i>Scopuloides hydnoides</i> *	<i>Scopuloides hydnoides</i>	KHL11916 (GB)	EU118665 <sup>c</sup>	EU118665 <sup>c</sup>
<i>Scopuloides hydnoides</i> *	<i>Scopuloides hydnoides</i>	GEL3859	AJ406573	–
<i>Scopuloides hydnoides</i> *	<i>Scopuloides hydnoides</i>	GEL3139	AJ406574	–
<i>Steccherinum ochraceum</i> *	<i>Steccherinum ochraceum</i>	KHL11902 (GB)	JQ031130	–
<i>Steccherinum ochraceum</i> *	<i>Steccherinum ochraceum</i>	Ryberg sn. (GB)	EU118670	–
Specimens “incertae sedis”	<i>Mycoacia aurea</i>	GEL5339	AJ406535	–
	<i>Mycoacia aurea</i>	NH14434	AY586691	–
	<i>Phlebia setulosa</i>	PH106520, culture	GU461311	–
	<i>Phlebia setulosa</i>	PH11749, culture	GU461312	–
	<i>Phlebia setulosa</i>	PH5105, culture	GU461313	–
	<i>Phlebia setulosa</i>	AH31879	GQ259417	–
	<i>Phlebia subochracea</i>	FCUG 1161, culture	AF141630	–
	<i>Phlebia subochracea</i>	KGN 162/95 (GB)	EU118656 <sup>b</sup>	EU118656 <sup>b</sup>
	<i>Phlebia suserialis</i>	FCUG1434, culture	AF141631	–
	<i>Phlebiella griseofulva</i>	GEL4492	AJ406517	–

<sup>a</sup>Names as indicated in Floudas & Hibbett (2015)

<sup>b</sup>Unpublished sequence

<sup>c</sup>These sequences contain part of SSU, complete ITS region (ITS1 + 5.8S + ITS2) and D1-D2 of LSU nrDNA.

EMBL/GenBank/DDBJ databases (Cochrane et al. 2011), mainly from Hibbett et al. (2000), Parmasto and Hallenberg (2000), Thorn et al. (2000), Hibbett and Binder (2002), Langer (2002), Larsson et al. (2004), Binder et al. (2005), Han et al. (2005), Kim et al. (2005), Larsson (2007), Hallenberg et al. (2008), Porter et al. (2008), Moreno et al. (2011), Miettinen et al. (2012), and Sjökvist et al. (2012), Floudas and Hibbett (2015). In order to clearly identify the genus of the Macaronesian specimens, we selected reference sequences from some of the genera included in Meruliaceae by



Larsson (2007), and from genera included by Wu et al. (2010) in his clade V that covers *Phlebia* sensu stricto and several taxa with odontoid or hydroid hymenophore. Moreover, sequences of type species of different genera listed in MycoBank (Crous et al. 2004, Robert et al. 2013, <http://www.mycobank.org>) as belonging to Meruliaceae were selected from EMBL/GenBank/DDBJ databases, mainly from references mentioned above (Table 1). Based on Binder et al. (2005) and Floudas and Hibbett (2015), two sequences of *Steccherinum* Gray (residual polypore clade) were included as outgroup. Where ambiguities in the alignment occurred, the alignment generating the fewest potentially informative characters was chosen (Baum et al. 1994). Alignment gaps were marked “–”, unresolved nucleotides and unknown sequences were indicated with “N”.

A maximum parsimony analysis (MP) was carried out; minimum length Fitch trees were constructed using heuristic searches with tree-bisection-reconnection (TBR) branch swapping, collapsing branches if maximum length was zero and with the MulTrees option on in PAUP\*4.0b10 (Swofford 2003), with a default setting to stop the analysis at 100 trees. Gaps were treated as missing data. Nonparametric bootstrap (MP-BS) support (Felsenstein 1985) for each clade, based on 10,000 replicates using the fast-step option, was tested. The consistency index, CI (Kluge and Farris 1969), retention index, RI, and rescaled consistency index, RC (Farris 1989) were obtained. The maximum likelihood (ML) analysis was done in PAUP\*Version 4.0b10, with the GTR+I+G model selected by this programme; for assessing branch support, 1000 non-parametric bootstrap replicates (ML-BS) were performed with the fast-step option. A third analysis was done by a Bayesian approach (Larget and Simon 1999, Huelsenbeck et al. 2001) using MrBayes 3.2 (Ronquist et al. 2012) and assuming the general time reversible model (Rodríguez et al. 1990), including estimation of invariant sites and a discrete gamma distribution with six categories (GTR+I+G), as selected by PAUP\*Version 4.0b10. Two independent and simultaneous analyses starting from different random trees were run for two million generations with 12 parallel chains, and trees and model scores saved every 100<sup>th</sup> generation. The initial 1000 trees were discarded as burn-in before calculating the 50% majority-rule consensus tree and the posterior probability (PP) of the nodes, as described in Telleria et al. (2010). A combination of bootstrap proportions and posterior probabilities was used to assess the level of confidence for a specific node (Lutzoni et al. 2004; Wilson et al. 2012). The phylogenetic trees were viewed with FigTree v. 1.3.1 (<http://tree.bio.ed.ac.uk/software/figtree/>) and edited with Adobe Illustrator CS3 v. 11.0.2 (Adobe Systems).

For molecular characterization of the Macaronesian specimens, the ITS sequences were compared with homologous sequences retrieved from the EMBL/GenBank/DDBJ (Cochrane et al. 2011, 2016) and UNITE (Abarenkov et al. 2011, Kõljag et al. 2013, <http://unite.ut.ee/cite.php>) databases (Table 1), mainly from de Koker et al. (2003), Larsson (2007), Hilden et al. (2008), Wu et al. (2010), Sjökvist et al. (2012), and Floudas and Hibbett (2015).

Based on our previous phylogenetic trees obtained from LSU, two sequences of *Phlebia radiata* Fr. were selected as outgroup (AY219366, EF491867). Alignment gaps were marked “–”, unresolved nucleotides and non-sequenced nucleotide positions within the data matrix were indicated with “N”. A maximum parsimony analysis (MP)



was carried out under heuristic search, following the same criteria as mentioned above for LSU; maximum likelihood (ML) and Bayesian approaches were also performed, using the GTR+I+G as selected by PAUP\*Version 4.0b10 and MrModeltest 2.3. The ML and Bayesian analyses were done with the same programs, and followed the same criteria as mentioned above for LSU.

Alignments and phylogenetic trees have been deposited at TreeBase: <http://purl.org/phylo/tree-base/phyloids/study/TB2:S21012>

## Results

Sixteen new sequences from the Macaronesian specimens were generated in this study (Table 1). The LSU sequence contains the domain D1-D2, and the ITS sequence the regions ITS1, 5.8S nrDNA and ITS2.

### LSU analyses


The LSU dataset contains 57 sequences and 908 aligned positions, of which 682 were constant, 82 parsimony uninformative, and 144 parsimony-informative. Maximum parsimony analysis yielded 100 most parsimonious trees (613 steps long, CI = 0.4731, HI = 0.6164, RI = 0.7399) under a heuristic search. Almost identical tree topologies were generated after parsimony and Bayesian analyses. The 50% majority-rule consensus tree from the Bayesian analysis is shown in Fig. 1, with percentage of bootstrap (MP-BS and ML-BS) and posterior probabilities indicated on the branches. The circumscription of clade V from Wu et al. (2010) is indicated in this figure.

All sequences obtained from Macaronesian specimens cluster in a supported clade (MP-BS = 70%, ML-BS = 80 %, PP = 1.0), with sequences from de Koker et al. (2003) and Floudas and Hibbett (2015) under *Hydnophlebia*. Most sequences are distributed over two supported clades, one containing four specimens from São Vicente (MP-BS = 96%, ML-BS = 95 %, PP = 1.0), and the other three specimens from Canary Islands (MP-BS = 93%, ML-BS = 95 %, PP = 0.98). Specimen 19071Tell., from Fogo Island, appears together with sequences KP135334 and KP135218, under *H. omnivora* 1 and *H. omnivora* 2 in Floudas and Hibbett (2015); although this relationship is not well supported (MP-BS = 93%, ML-BS = 95 %, PP = 0.98). The two *H. chrysorhizon* sequences (AF139967 and KP135217 from previous authors) form a fairly well supported clade with the specimens from São Vicente (MP-BS = 61%, ML-BS = 65 %, PP = 1.0).

### ITS analyses

The ITS nrDNA dataset contains 26 sequences and 851 aligned positions, of which 575 were constant, 103 parsimony uninformative, and 173 parsimony-informative. After heuristic search, the 100 trees had 447 steps, CI = 0.7136, HI = 0.3798 and RI






**Figure 1.** The 50% majority rule Bayesian tree inferred from D1-D2 LSU nrDNA assuming the GTR + I + G model of corticioid fungi included in Table 1. Parsimony bootstrap values (> 50%) maximum likelihood bootstrap values (> 50%) and Bayesian posterior probabilities (> 0.95) are indicated on the branches. Clade V from Wu et al. (2010) is indicated. Taxon name between parentheses indicate specimens with uncertain generic placement. Sequences of the new species described in this paper, *H. canariensis*, *H. gorgonea* and *H. meloi* are in bold. The asterisk (\*) after the taxon names denotes type species of the genus.

= 0.7831. Almost identical tree topologies were generated after parsimony (data not shown), maximum likelihood (data not shown) and Bayesian analyses. The 50% majority-rule consensus tree from the Bayesian analysis is shown in Fig. 2, with percentage of bootstrap (MP-BS and ML-BS), and posterior probabilities indicated on the branches.





**Figure 2.** The 50% majority rule Bayesian tree inferred from ITS nrDNA assuming the GTR + I + G model of corticioid fungi included in Table 1. Parsimony bootstrap values (> 50%), maximum likelihood bootstrap values (> 50%) and Bayesian posterior probabilities (> 0.95) are indicated on the branches. Taxon name between parentheses indicate specimens with uncertain generic placement. Sequences of the new species described in this paper, *H. canariensis*, *H. gorgonea* and *H. meloi* are in bold. The asterisk (\*) after the taxon names denotes type species of the genus.

Similar to the LSU analyses, the sequences from Macaronesian specimens form a clade (MP-BS = 76%, ML-BS = 75%, PP = 1.0), together with downloaded sequences of *Hydnophlebia* from the USA (Arizona, Florida, Illinois, New York and Puerto Rico) and Canada, identified in Floudas and Hibbett (2015) as *H. chrysorhizon*, *H. cf. chrysorhizon*, *H. omnivora* 1 and *H. omnivora* 2; as well as sequences of *H. chrysorhizon* from Illinois and *H. omnivora* from Arizona published by de Koker et al. (2003) that we consider represent the modern interpretation of these two species. Sequence UDB016816 from Madagascar, in Fig. 2, labelled as *Hydnophlebia* sp. 2, (under *Phlebia* sp. in UNITE database) also clusters in this clade.

The five sequences from Canada and the USA identified as *H. chrysorhizon* grouped in a highly supported clade (MP-BS > 80%, ML-BS > 82%, PP = 1.0). The sequences



UDB016816, labelled *Hydnophlebia* sp. 2 from Madagascar, and KP135339, labelled *Hydnophlebia* sp. 3 (*H. cf. chrysorhizon* in Floudas and Hibbett 2015) from Puerto Rico, did not group with other *Hydnophlebia* sequences.

The new sequences generated for this work are distributed over three clades. These clades are here described in the order they occur from top to bottom in Fig. 2.

The first group (MP-BS = 100%, ML-BS = 100%, PP = 1.0) contains sequences of 17035Tell., 17038Tell., and 17674Tell. from El Hierro and Fuerteventura Islands, collected on different substrates.

A second clade (MP-BS > 75%, ML-BS > 82%, PP = 1.0) encloses specimen 19071Tell., from Fogo Island, collected on *Sarcostemma daltonii* Decne (Asclepiadaceae), and the sequence indicated in Fig. 2 as *Hydnophlebia* sp. 1 (*H. omnivora* 1 in Floudas and Hibbett 2015) and three sequences of *Hydnophlebia omnivora* (two of them as *H. omnivora* 2 in Floudas and Hibbett 2015).

Sequences of 13327MD, 19110Tell., 19111Tell., and 19133Tell., all from São Vicente Island on *Prosopis juliflora* (Sw.) DC. (Fabaceae), are distributed in a third clade (MP-BS = 100%, ML-BS = 99%, PP = 1.0).

## Taxonomy

***Hydnophlebia* Parmasto, Izv. Akad. Nauk Estonsk. SSR, Ser. Biol. 16: 384. 1967**

**Type species.** *Hydnum chrysorhizon* Torr. in Eaton, Manual Bot.: 309. 1822

Basidioma resupinate, separable, generally reddish orange yellow. Hymenophore hydroid, with aculei up to 0.5–1.5 mm long, conical to cylindrical. Margin with strands. Hyphal system monomitic with scattered clamps, subicular and strand hyphae thick-walled, colorless to yellowish, encrusted, aculei and subhymenial hyphae thin-walled, also colorless. Cystidia cylindrical to ventricose, colorless, thin-walled, sometimes few. Basidia cylindrical to subclavate, with 4 sterigmata, basal clamp absent. Spores cylindrical, ellipsoid to subglobose, smooth, thin-walled.

## Key to species of *Hydnophlebia*

- 1 Spores narrowly ellipsoid to cylindrical,  $4\text{--}6 \times 2\text{--}3 \mu\text{m}$  ( $L/W = 1.9$ ). Basidiome orange brown in dry specimens, reddish orange to deep orange in fresh material. Strands very long, yellowish to cream in dry material and reddish orange in fresh specimens, up to 1 mm diam ..... **2. *H. chrysorhizon***
- Spores ellipsoid, broadly ellipsoid or subglobose ( $L/W \leq 1.6$ ) ..... **2**
- 2 Spores subglobose,  $4\text{--}5.5 \times 3\text{--}4 \mu\text{m}$  ( $L/W = 1.2$ ). Basidiome yellowish white to pale orange-yellow in dry specimens. Margin fimbriate, yellowish white, with strands poorly developed. Cystidia cylindrical,  $40\text{--}55 \times 3\text{--}4 \mu\text{m}$  ..... **4. *H. meloi***
- Spores ellipsoid to broadly ellipsoid ( $L/W \geq 1.3$ ) ..... **3**



- 3 Basidiome in small and poorly developed patches, cream-coloured in dry specimens. Clamps present in strand hyphae. Cystidia cylindrical slightly tapered at the apex,  $40\text{--}70 \times 4\text{--}5 \mu\text{m}$ . Spores ellipsoid,  $5\text{--}6.5 \times 3\text{--}4 \mu\text{m}$  ( $L/W = 1.6$ ) ..... **5. *H. omnivora***
- Basidiome broadly effuse, orange-yellow to brilliant orange-yellow. Clamps absent in strand hyphae ..... **4**
- 4 Basidiome light orange-yellow in dry specimens. Margin fimbriate, with white, well developed strands. Cystidia of two types, cylindrical with slightly tapered apex and ventricose with subulate apex,  $45\text{--}55 \times 3\text{--}5 \mu\text{m}$ . Spores ellipsoid,  $5\text{--}7 \times 3\text{--}4.5 \mu\text{m}$  ( $L/W = 1.5$ ) ..... **1. *H. canariensis***
- Basidiome light to brilliant orange-yellow in dry specimens. Margin fimbriate with poorly developed strands. Cystidia cylindrical, sometimes capitate, to ventricose,  $45\text{--}55 \times 4\text{--}6 \mu\text{m}$ . Spores ellipsoid to broadly ellipsoid,  $5\text{--}7 \times 4\text{--}4.5 \mu\text{m}$  ( $L/W = 1.4$ ) ..... **3. *H. gorgonea***

**1. *Hydnophlebia canariensis* Telleria, M. Dueñas & M.P. Martín, sp. nov.**

MycoBank MB815729

Figs 3, 4

**Diagnosis.** This species can be recognized by the orange-yellow basidiome, hydroid hymenophore with long aculei, up to 1.5 mm, white subiculum, and well-developed white strands. Spores ellipsoid  $5\text{--}7 \times 3\text{--}4.5 \mu\text{m}$  ( $L/W = 1.5$ ).

**Type.** SPAIN. Canary Islands: El Hierro, Frontera, Sabinar de la Dehesa,  $27^{\circ}44'43''\text{N}$ ;  $18^{\circ}07'02''\text{W}$ , 610 m alt., on unidentified wood, 26 January 2007, M.T. Telleria, 17035Tell. (holotype: MA-Fungi 86622). LSU sequence KF528103, ITS sequence KF483012.

**Etymology.** Named after the Canary Islands where the holotype and paratypes were collected.

**Description.** Basidiome resupinate, effuse, membranous to ceraceous, yellow (82. v. Y) in fresh specimens and light orange-yellow (70. l. OY) in dry. Hymenophore hydroid, aculei conical, 0.5–1.5 mm long. Subiculum byssoid, white. Margin fimbriate, white, with well-developed white strands.

Hyphal system monomitic; subicular hyphae  $6\text{--}8 \mu\text{m}$  wide, with clamps, thin to thick-walled; strand hyphae  $7\text{--}11 \mu\text{m}$  wide, without clamps, thick-walled; aculei hyphae  $4\text{--}5 \mu\text{m}$  wide, without clamps, thin-walled and growing perpendicular to the substrate; subhymenial hyphae  $3\text{--}4 \mu\text{m}$  wide, without clamps, thin-walled, and loosely interwoven. Cystidia of two types: cylindrical with slightly tapered apex and ventricose with subulate apex, thin-walled,  $45\text{--}55 \times 3\text{--}5 \mu\text{m}$ . Basidia cylindrical to subclavate,  $24\text{--}28 \times 4\text{--}6 \mu\text{m}$ , with 4 sterigmata, basal clamp absent. Spores ellipsoid  $5\text{--}7 \times 3\text{--}4.5 \mu\text{m}$  ( $L/W = 1.5$ ), thin-walled, colorless, smooth.

**Ecology and distribution.** On decayed wood and plant debris in arid and semi-arid habitats; known only from the Canary Islands.






**Figure 3.** *Hydnophlebia canariensis*. **a, b** Collection 17035Tell., MA-Fungi 86622, holotype, basidiome, wet (a) and hymenophore, dry specimen (b) **c** Collection 17038Tell. MA-Fungi 86623, basidiome, dry specimen. *Hydnophlebia chrysorhizon*. Collection NY, lectotype **d** Basidiome, dry specimen **e** Hymenophore, dry specimen **f** Strands, dry specimen. Scale bars: **a, e** = 5 mm; **b** = 1.5 mm; **c, d, f** = 1 cm.

**Other specimens examined.** Spain. Canary Islands: El Hierro, Frontera, Sabinar de la Dehesa, 27°44'43"N; 18°07'02"W, 610 m alt., on unidentified wood, 26 January 2007, M.T. Telleria, 17038Tell. (MA-Fungi 86623), LSU sequence KF528104, ITS sequence KF483013. Fuerteventura, Pájara, Parque Natural de Jandía, Valle de los Mosquitos, 28°04'36"N; 14°25'23"W, 99 m alt., on *Launaea arborescens*, 05 December 2007, M.T. Telleria, 17674Tell. (MA-Fungi 86619), LSU sequence KF528100, ITS sequence KF483009.





**Figure 4.** *Hydnophlebia canariensis*. Collection 17035 Tell., MA-Fungi 86622, holotype **a** Hymenophore **b** Vertical section through an aculei **c** Subicular hyphae **d** Strand hyphae **e** Aculei hyphae **f** Subhymenial hyphae, cystidia, and basidia **g** Spores. Scale bars: **a** = 1 mm; **b** = 25  $\mu$ m; **c–g** = 10  $\mu$ m. Drawing by M. Dueñas.



**2. *Hydnophlebia chrysorhizon* (Torr.) Parmasto, Izv. Akad. Nauk Estonsk. SSR, Ser. Biol. 16: 384. 1967**

Figs 3, 5

**Basionym.** *Hydnum chrysorhizon* Torr. in Eaton, Manual Bot.: 309. 1822

**Type.** USA, *Hydnum chrysorhizon* Torr. in Eaton Man. 3ed. p. 309. 237, C. C., Steward. In herbarium NY! (lectotype, designated by Burdsall 1985).

**Description.** Basidiome resupinate, effuse, membranous, easily separable, orange-brown in dry specimens, reddish orange to deep orange in fresh material (Burdsall and Nakasone 1978, Lindsey and Gilbertson 1975, Burdsall 1985, Maekawa 1993). Hymenophore hydroid, aculei dense, conical to subcylindrical, 1–1.6 mm long. Margin with strands very long and well developed, yellowish to cream in dry specimens, reddish orange in fresh specimens (Burdsall 1985), up to 1 mm diam.

Hyphal system monomitic; subicular hyphae 7–10 µm wide, with clamps, thick-walled, colorless to pale yellow, densely encrusted with colorless crystals and loosely interwoven; strand hyphae 10–17 µm wide, without clamps, thick-walled, colorless, also encrusted; aculei hyphae 4–6 µm wide, with scattered clamps, thin-walled, colorless, and oriented perpendicular to the substrate; subhymenial hyphae 5–7 µm wide, without clamps, thin-walled, colorless, densely interwoven, short-celled. Cystidia not seen, but according to Burdsall (1985) few, cylindrical, thin-walled, hyaline, short, 18–40 × 4.5–6 µm. Basidia clavate, 15–21 × 4–6 µm, with 4 sterigmata, basal clamp absent. Spores narrowly ellipsoid to cylindrical, 4–6 × 2–3 µm (L/W = 1.9), thin-walled, colorless, smooth.


**Ecology and distribution.** On decayed wood. Described from New York (Eaton 1822), this species has been reported from: Africa: Cameroon, (Roberts 2000), and Seychelles (Hjortstam and Ryvarden 2009); North America: USA, Arizona, Florida, Maryland, Mississippi, New York, North Carolina, South Carolina, Tennessee, Wisconsin (Lindsey and Gilbertson 1975, Burdsall and Nakasone 1978, Burdsall 1985, Nakasone 2012); South America: Argentina, Venezuela, Brazil (Hjortstam and Ryvarden 2007); Meso America: Puerto Rico (Hjortstam and Ryvarden 2009), as well as Saint Vincent and the Grenadines (Nakasone 2012); Asia: Japan (Maekawa 1993).

**Other specimens examined.** USA. Ohio, Hamilton Co. Sharon Woods County Park, on *Quercus* sticks, 13 October 1973, W.B. & V.G. Cooke 48958. New York, New Dorp, Staten Island, 17 October 1896, col. L.M. Underwood.

**Remarks.** This species has very long and well-developed strands and, microscopically, it is the only species in the genus with spores narrowly ellipsoid to cylindrical (L/W = 1.9) and scattered clamps in the aculei hyphae.

Based on morphological analyses, Burdsall (1985) considered *Hydnum fragilissimum* Berk. & M.A. Curtis, *Hydnum ischnodes* Berk., and *Hydnum chrysocomum* Underw. as synonyms of *H. chrysorhizon*; and according to Nakasone (2012) *Hydnum schweinitzii* Berk. & M.A. Curtis, *Hydnum chrysodon* Berk. & M.A. Curtis, and *Merulius elliotii* Masee are other synonyms.





**Figure 5.** *Hydnophlebia chrysorhizon*. Collection NY, lectotype **a** Hymenophore **b** Vertical section through an aculei **c** Subicular hyphae **d** Strand hyphae **e** Aculei hyphae **f** Subhymenial hyphae and basidia **g** Spores. Scale bars: **a** = 1mm; **b** = 25  $\mu$ m; **c–g** = 10  $\mu$ m. Drawing by M. Dueñas.



**3. *Hydnophlebia gorgonea* Telleria, M. Dueñas & M.P. Martín, sp. nov.**

MycoBank MB815730

Figs 6, 7

**Diagnosis.** Morphologically this species is similar to *Hydnophlebia canariensis*, but can be distinguished by the strands, well developed in *H. canariensis* and poorly so in *H. gorgonea*. Spores ellipsoid to broadly ellipsoid  $5\text{--}7 \times 4\text{--}4.5 \mu\text{m}$  ( $L/W = 1.4$ ).

**Type.** CAPE VERDE. São Vicente: Mindelo, Ribeira da Vinha,  $16^{\circ}51'49''\text{N}$ ;  $25^{\circ}00'09''\text{W}$ , 10 m alt., on *Phoenix atlantica*, 26 September 2010, M.T. Telleria, 19111Tell. (holotype: MA-Fungi 86659), LSU sequence KF528140, ITS sequence KF483049.

**Etymology.** Named after Gorgades, an ancient name for the Cape Verde Islands, Atlantic Ocean.

**Description.** Basidiome resupinate, effuse, membranous, easily separable, light orange-yellow (70. l. OY) to brilliant orange-yellow (67. brill. OY). Hymenophore hydroid, aculei conical, 0.6–1 mm long. Margin fimbriate, white, with poorly developed strands.

Hyphal system monomitic; subicular hyphae 6–8  $\mu\text{m}$  wide, with clamps, thin- to thick-walled, loosely interwoven, hyaline, encrusted with colorless crystals; strand hyphae 12–15  $\mu\text{m}$  wide, without clamps, thick-walled, sometimes gelatinous and also encrusted; aculei hyphae 3.5–4.5  $\mu\text{m}$  wide, without clamps, thin-walled, colorless, growing perpendicular to the substrate; subhymenial hyphae 4.5–8  $\mu\text{m}$  wide, without clamps, thin-walled, colorless, loosely interwoven, and short- to long-celled. Cystidia cylindrical to ventricose, sometimes capitate, thin-walled,  $45\text{--}55 \times 4\text{--}6 \mu\text{m}$ . Basidia cylindrical to subclavate,  $22\text{--}24 \times 6\text{--}8 \mu\text{m}$ , with 4 sterigmata, basal clamp absent. Spores ellipsoid to broadly ellipsoid  $5\text{--}7 \times 4\text{--}4.5 \mu\text{m}$  ( $L/W = 1.4$ ), thin-walled, colorless, smooth.

**Ecology and distribution.** This species is known from only two localities of São Vicente Island, Cape Verde Archipelago, on decayed wood of *Phoenix atlantica* and *Prosopis juliflora* in arid habitats.

**Other specimens examined.** CAPE VERDE. São Vicente: Mindelo, Ribeira da Vinha,  $16^{\circ}51'49''\text{N}$   $25^{\circ}00'09''\text{W}$ , 10 m alt., on *Prosopis juliflora*, 26 September 2010, M.T. Telleria, 19110Tell. (MA-Fungi 86658), LSU sequence KF528139, ITS sequence KF483048; M. Dueñas, 13327MD (MA-Fungi 86642), LSU sequence KF528122, ITS sequence KF483031. São Vicente: Ermida,  $16^{\circ}50'26''\text{N}$ ;  $24^{\circ}57'23''\text{W}$ , 100 m alt., on *Prosopis juliflora*, 26 September 2010, M.T. Telleria, 19133Tell. (MA-Fungi 86664), LSU sequence KF528145, ITS sequence KF483054.


**4. *Hydnophlebia meloi* Telleria, M. Dueñas & M.P. Martín, sp. nov.**

MycoBank MB815731

Figs 6, 8

**Diagnosis.** Similar to *Hydnophlebia omnivora* but differs in having subglobose spores,  $4\text{--}5.5 \times 3\text{--}4 \mu\text{m}$  ( $L/W = 1.2$ ), instead of ellipsoid,  $5\text{--}6.5 \times 3\text{--}4 \mu\text{m}$  ( $L/W = 1.6$ ). This is the only species in the genus with subglobose spores.






**Figure 6.** *Hydnophlebia gorgonea*. **a** Collection 19111Tell., MA-Fungi 86659, holotype, basidiome, dry specimen **b** Collection 19133Tell., MA-Fungi 86664, basidiome, dry specimen. *Hydnophlebia meloi* **c**, **d** Collection 19071Tell., MA-Fungi 86654, holotype, basidiome, dry specimen (**c**), and margin and strands, dry specimen (**d**). *Hydnophlebia omnivora* **e** Collection 5267 C.R. Shear coll., BPI, holotype **f** Basidiome, dry specimen. Scale bars: **a–b** = 5 mm; **c–d**, **f** = 2 mm; **e** = 10 mm.

**Type.** CAPE VERDE. Fogo: Mosteiros, Miradouro, 15°01'41"N; 24°19'13"W, 283 m alt., on *Sarcostemma daltonii*, 24 September 2010, M.T. Telleria, 19071Tell. (holotype: MA-Fungi 86654), LSU sequence KF528135, ITS sequence KF483044.

**Etymology.** Named after Ireneia Melo, colleague and friend, Portuguese mycologist from the Botanical Garden of the University of Lisbon.






**Figure 7.** *Hydnophlebia gorgonea*. Collection 19111Tell., MA-Fungi 86659, holotype **a** Hymenophore **b** Vertical section through an aculei **c** Subicular hyphae **d** Strand hyphae **e** Aculei hyphae **f** Subhymenial hyphae, cystidia, and basidia **g** Spores. Scale bars: **a** = 1 mm; **b** = 25  $\mu$ m; **c–g** = 10  $\mu$ m. Drawing by M. Dueñas.

**Description.** Basidiome resupinate, effuse, membranous to ceraceous, yellowish white (92. y White) to pale orange-yellow (73. p. OY) in dry specimens. Hymenophore hydroid, aculei conical, 0.5–1 mm long, in dried specimens usually broken. Margin fimbriate, yellowish white, with strands.





**Figure 8.** *Hydnophlebia meloi*. Collection 19071Tell., MA-Fungi 86654, holotype **a** Hymenophore **b** Vertical section through an aculei **c** Subicular hyphae **d** Strand hyphae **e** Aculei hyphae **f** Subhymenial hyphae, cystidia, and basidia **g** spores. Scale bars: **a** = 1 mm; **b** = 25  $\mu$ m; **c–g** = 10  $\mu$ m. Drawing by M. Dueñas.

Hyphal system monomitic; subicular hyphae 6–7.5  $\mu$ m wide, with clamps, thick-walled, loosely interwoven; strand hyphae 6–7.5  $\mu$ m wide, with clamps occasionally double, thick-walled, sometimes encrusted with colorless crystals; aculei hyphae 3.5–4.5  $\mu$ m wide, without clamps, thin-walled, growing perpendicular to the substrate; sub-



hymenial hyphae 3.5–4.5  $\mu\text{m}$  wide, without clamps, thin-walled, loosely interwoven, short- to long-celled. Cystidia cylindrical, thin-walled, 40–55  $\times$  3–4  $\mu\text{m}$ . Basidia cylindrical to subclavate, 18–26  $\times$  5–7  $\mu\text{m}$ , with 4 sterigmata, basal clamp absent. Spores subglobose 4–5.5  $\times$  3–4  $\mu\text{m}$  (L/W = 1.2), thin-walled, colorless, smooth.

**Distribution.** Rocky steep slopes, on *Sarcostemma daltonii*, endemic climbing herb of Cape Verde Archipelago. Only known from the type locality in Fogo Island.

**Other specimens examined.** CAPE VERDE. Fogo: Mosteiros, Miradouro, 15°01'41"N; 24°19'13"W, 283 m alt., on *Sarcostemma daltonii*, 24 September 2010, M.T. Telleria, 19072Tell. (MA-Fungi 90746).

## 5. *Hydnophlebia omnivora* (Shear) Hjortstam & Ryvarden, Synopsis Fungorum 26: 10-23. 2009

Figs 6, 9

**Basionym.** *Hydnum omnivorum* Shear, J. Agric. Res. 30: 476. 1925

**Type.** USA, C.L.S. Type on Osage Orange [*Macura pomifera*], near Paris, Texas. C.R. Shear coll. Sept. 1903, no. 5267. In herbarium BPI! (holotype).


**Description.** Basidiome effuse in small and poorly developed patches, cream-coloured in dry specimens. Hymenophore, according to Burdsall (1985), hydroid, aculei conical to subcylindrical, 0.6–1 mm long; broken or poorly developed in type material. Margin floccose to fibrillose, white, with strands poorly developed.

Hyphal system monomitic; subicular hyphae 8–11  $\mu\text{m}$  wide, with clamps occasionally double, thick-walled, loosely interwoven; strand hyphae 5–9  $\mu\text{m}$  wide, with a few clamps, thick-walled, colorless; aculei hyphae 4–5  $\mu\text{m}$  wide, without clamps, thin-walled, growing perpendicular to the substrate; subhymenial hyphae 5–6  $\mu\text{m}$  wide, without clamps, thin-walled, densely interwoven, short-celled. Cystidia cylindrical, slightly tapered to apex, thin-walled, 40–70  $\times$  4–5  $\mu\text{m}$ . Basidia cylindrical to subclavate, 17–21  $\times$  6–7  $\mu\text{m}$ , with 4 sterigmata, basal clamp absent. Spores ellipsoid, 5–6.5  $\times$  3–4  $\mu\text{m}$  (L/W = 1.6), thin-walled, colorless, smooth.

**Ecology and distribution.** Described from Texas (Shear 1925). According to Burdsall (1985) this species is distributed in the arid regions of southwestern United States, and probably into southern California and northern Mexico. Also reported from Florida (Ginns and Lefebvre 1993) and Uruguay (Martínez and Nakasone 2005).

**Remarks.** Molecular analyses indicate that this species is related to *H. meloi*. Morphologically they can be distinguished by the shape and size of spores, subglobose 4–5.5  $\times$  3–4  $\mu\text{m}$  in *H. meloi*, and ellipsoid 5–6.5  $\times$  3–4  $\mu\text{m}$  in *H. omnivora*.





**Figure 9.** *Hydnophlebia omnivora*. Collection 5267 C.R. Shear coll., BPI, holotype **a** Hymenophore **b** Vertical section through an aculei **c** Subicular hyphae **d** Strand hyphae **e** Aculei hyphae **f** Subhymenial hyphae, cystidia, and basidia **g** Spores. Scale bars: **a** = 1 mm; **b** = 25  $\mu$ m; **c–g** = 10  $\mu$ m. Drawing by M. Dueñas.



## Discussion

In this study a taxonomic analysis of *Hydnophlebia*, based on morphological and molecular data, is provided. *Hydnophlebia* has been confused with *Phanerochaete* and the two species included, *Hydnophlebia chrysorhizon* and *Hydnophlebia omnivora*, were assigned to the latter genus (Burdall 1985).

For a long time, *Hydnophlebia* was considered a monospecific genus; however, based on the molecular analyses, both LSU and ITS sequences, as well as a point-by-point comparison of the morphological characters, five species can be discriminated, two already described by other authors (*H. chrysorhizon* and *H. omnivora*), and the three new species from Macaronesia described here (*H. canariensis*, *H. gorgonea*, and *H. meloi*).

Moreover our results show that three other species could be described, although more collections should be analysed: 1) *Hydnophlebia* sp. 1 under *H. omnivora* 1 in Floudas and Hibbett (2015); 2) *Hydnophlebia* sp. 2 under *Phlebia* sp. from Madagascar in UNITE database; and 3) *Hydnophlebia* sp. 3 under *Hydnophlebia* cf. *chrysorhizon* in Floudas and Hibbett (2015).

## Acknowledgements

Financial support was provided by DGICT projects CGL2012-35559, CGL2015-67459-P. We are grateful to reviewers for comments and suggestions to improve the final version. Also to Marian Glenn for checking the English, to Fátima Durán (RJB/CSIC) for providing technical assistance, and to the curators of BPI and NY herbaria for their invaluable assistance.

## References

- Abarenkov K, Nilsson RH, Larsson KH, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjøller R, Larsson E, Pennanen T, Sen R, Taylor AF, Tedersoo L, Ursing BM, Vrålstad T, Liimatainen K, Peintner U, Kõljalg U (2011) The UNITE database for molecular identification of fungi – recent updates and future perspectives. *New Phytologist* 186: 281–285. <https://doi.org/10.1111/j.1469-8137.2009.03160.x>
- Altschul SE, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Research* 25: 3389–3402. <https://doi.org/10.1093/nar/25.17.3389>
- Baum DA, Sytsma K, Hoch P (1994) A molecular phylogenetic analysis of *Epilobium* based on sequences of nuclear ribosomal DNA. *Systematic Botany* 19: 363–388. <https://doi.org/10.2307/2419763>
- Binder M, Hibbett DS, Larsson KH, Langer E, Langer G (2005) The phylogenetic distribution of resupinate forms across the major clades of homobasidiomycetes. *Systematics and Biodiversity* 3: 113–157. <https://doi.org/10.1017/S1477200005001623>



- Budington AB, Gilbertson RL (1973) Some Southwestern lignicolous Hymenomycetes of special interest. *Southwestern Naturalist* 17: 409–422. <https://doi.org/10.2307/3670128>
- Burdsall HH (1985) A contribution to Taxonomy of the Genus *Phanerochaete*. *Mycologia Memoir* 10: 1–165.
- Burdsall HH, Nakasone KK (1978) Taxonomy of *Phanerochaete chrysorhizon* and *Hydnum omnivorum*. *Mycotaxon* 7: 10–22.
- Cochrane G, Karsch-Mizrachi I, Nakamura Y (2011) The International Nucleotide Sequence Database Collaboration. *Nucleic Acids Research* 39: 15–18. <https://doi.org/10.1093/nar/gkq1150>
- Cochrane G, Karsch-Mizrachi I, Nakamura Y (2016) The International Nucleotide Sequence Database Collaboration. *Nucleic Acids Research* 44: 48–50. <https://doi.org/10.1093/nar/gkv1323>
- Crous PW, Gams W, Stalpers JA, Robert V, Stegehuis G (2004) MycoBank: an online initiative to launch mycology into the 21<sup>st</sup> century. *Studies in Mycology* 50: 19–22.
- de Koker TH, Nakasone KK, Haarhof J, Burdsall HH Jr, Janse BJ (2003) Phylogenetic relationships of the genus *Phanerochaete* inferred from the internal transcribed spacer region. *Mycological Research* 107: 1032–1040.
- Eaton M (1822) *Manual of Botany for the Northern and Middle States of America*. Albany.
- Farris JS (1989) The retention index and the rescaled consistency index. *Cladistics* 5: 417–419. <https://doi.org/10.1111/j.1096-0031.1989.tb00573.x>
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
- Floudas D, Hibbett DS (2015) Revisiting the taxonomy of *Phanerochaete* (Polyporales, Basidiomycota) using a four gene dataset and extensive ITS sampling. *Fungal Biology* 119: 679–719. <https://doi.org/10.1016/j.funbio.2015.04.003>
- Gardes M, Bruns TD (1993) ITS Primers with enhanced specificity for Basidiomycetes-applications to the identification of mycorrhizae and rusts. *Molecular Ecology* 1: 113–118. <https://doi.org/10.1111/j.13625-294X.1993.tb00005.x>
- Ginns J, Lefebvre MNL (1993) Lignicolous corticioid fungi (Basidiomycota) of North America. *Mycologia Memoir* 19: 1–247.
- Hallenberg N, Ryberg M, Nilsson RH, Wood AR, Wu SH (2008) *Pseudolagarobasidium* (Basidiomycota): the reinstatement of a genus of parasitic, saprophytic, and endophytic resupinate fungi. *Botany* 86: 1319–1325. <https://doi.org/10.1139/B08-088>
- Han M-J, Choi H-T, Song H-G (2005) Purification and characterization of Laccase from the white root fungus *Trametes versicolor*. *The Journal of Microbiology* 43: 555–560.
- Hibbett DS, Binder M (2002) Evolution of complex fruiting-body morphologies in homobasidiomycetes. *Proceedings of the Royal Society of London, Series B, Biological Sciences* 269: 1963–1969. <https://doi.org/10.1098/rspb.2002.2123>
- Hibbett DS, Gilbert LB, Donoghue MJ (2000) Evolutionary instability of ectomycorrhizal symbioses in basidiomycetes. *Nature* 407: 506–508. <https://doi.org/10.1038/35035065>
- Hilden KS, Bortfeldt R, Hofrichter M, Hatakka A, Lundell TK (2008) Molecular characterization of the basidiomycete isolate *Nematoloma frowardii* b19 and its manganese peroxidase places the fungus in the corticioid genus *Phlebia*. *Microbiology*. 154: 2371–2379. <https://doi.org/10.1099/mic.0.2008/018747-0>



- Hjortstam K, Ryvarden L (2007) Checklist of corticioid fungi (Basidiomycotina) from the tropics, subtropics and the southern hemisphere. *Synopsis Fungorum* 22: 27–146.
- Hjortstam K, Ryvarden L (2009) A preliminary checklist of Aphyllophorales from the Seychelles. *Synopsis Fungorum* 26: 10–23.
- Huelsenbeck JP, Ronquist F, Nielsen R, Bollback JP (2001) Bayesian inference of phylogeny and its impact on evolutionary biology. *Science* 294: 2310–2314. <https://doi.org/10.1126/science.1065889>
- Kelly KL, Judd DB (1976) COLOR. Universal language and dictionary of names. National Bureau of Standards, Special Publication 440, 184 pp.
- Kim G-H, Lim YW, Song Y-S, Kim J-J (2005) Decay fungi from playground wood products in service using 28S rDNA sequence analysis. *Holzforschung* 59: 459–466. <https://doi.org/10.1515/HF.2005.076>
- Kluge AG, Farris JS (1969) Quantitative phyletics and the evolution of anurans. *Systematic Zoology* 18: 1–32. <https://doi.org/10.2307/2412407>
- Kõljalg U, Nilsson RH, Abarenkov K, Tedersoo L, Taylor AF, Bahram M, Bates ST, Bruns TD, Bengtsson-Palme J, Callaghan TM, Douglas B (2013) Towards a unified paradigm for sequence-based identification of Fungi. *Molecular Ecology* 22: 5271–5277. <https://doi.org/10.1111/mec.12481>
- Langer E (2002) Phylogeny of non-gilled and gilled basidiomycetes: DNA sequence inference, ultrastructure and comparative morphology. Habilitationsschrift, Tübingen University, Tübingen, Germany.
- Larget B, Simon DL (1999) Markov chain Monte Carlo algorithms for the Bayesian analysis of phylogenetic trees. *Molecular Biology and Evolution* 16: 750–759. <https://doi.org/10.1093/oxfordjournals.molbev.a026160>
- Larsson KH (2007) Re-thinking the classification of corticioid fungi. *Mycological Research* 111: 1040–1063. <https://doi.org/10.1016/j.mycres.2007.08.001>
- Larsson KH, Larsson E, Kõljalg U (2004) High phylogenetic diversity among corticioid homobasidiomycetes. *Mycological Research* 108: 983–1002. <https://doi.org/10.1017/S0953756204000851>
- Lindsey JP, Gilbertson RL (1975) Wood-inhabiting Homobasidiomycetes on Saguaro in Arizona. *Mycotaxon* 2: 83–103.
- Lutzoni F, Kauff F, Cox CJ, McLaughlin D, Celio G, Dentinger B, Padamsee M, Hibbett D, James TY, Baloch E, Grube M (2004) Assembling the fungal tree of life: progress, classification, and evolution of subcellular traits. *American Journal of Botany* 91: 1446–1480. <https://doi.org/10.3732/ajb.91.10.1446>
- Maekawa N (1993) Taxonomic study of Japanese Corticiaceae (Aphyllophorales). I. Report Tottori Mycological Institute 31: 1–149.
- Martín MP, Winka K (2000) Alternative methods of extracting and amplifying DNA from lichens. *Lichenologist* 32: 189–196. <https://doi.org/10.1006/lich.1999.0254>
- Martínez S, Nakasone K (2005) The genus *Phanerochaete* (Corticiaceae, Basidiomycotina) sensu lato in Uruguay. *Sydowia* 57: 94–101.
- Miettinen O, Larsson E, Sjökvist E, Larsson KH (2012) Comprehensive taxon sampling reveals unaccounted diversity and morphological plasticity in a group of dimitic polypores



- (Polyporales, Basidiomycota). *Cladistics* 28: 251–270. <https://doi.org/10.1111/J.1096-0031.2011.00380.X>
- Moreno G, Blanco M-N, Checa J, Plata G, Peláez F (2011) Taxonomic and phylogenetic revision of three rare irpicoid species within the Meruliaceae. *Mycological Progress* 10: 481–491. <https://doi.org/10.1007/s11557-010-0717-y>
- Nakasone KK (2012) Type studies of corticioid Hymenomycetes (Basidiomycota) with aculei – Part II. *Czech Mycology* 64: 23–42.
- Parmasto E (1967) Corticiaceae URSS IV. Descriptiones taxorum novarum. Combinationes novae. *Eesti NSV Teaduste Akadeemia Toimetised* 16: 377–394.
- Parmasto E (1968) *Conspectus Systematis Corticiacearum*. Institutum Zoologicum et Botanicum Academiae Scientiarum Estonicae, Tartu.
- Parmasto E, Hallenberg N (2000) A taxonomic study of Phlebioid fungi (Basidiomycota). *Nordic Journal of Botany* 20: 105–118. <https://doi.org/10.1111/j.1756-1051.2000.tb00740.x>
- Porter TM, Skillman JE, Moncalvo JM (2008) Fruiting body and soil rDNA sampling detects complementary assemblage of Agaricomycotina (Basidiomycota, Fungi) in a hemlock-dominated forest plot in southern Ontario. *Molecular Ecology* 17: 3037–3050. <https://doi.org/10.1111/j.1365-294X.2008.03813.x>
- Rambaut A (2002) Se-Al: sequence alignment editor v2.0a11. Institute of Evolutionary Biology, University of Edinburgh.
- Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analyzed from nuclear large subunit ribosomal DNA sequences. *Mycological Research* 98: 625–634. [https://doi.org/10.1016/S0953-7562\(09\)80409-7](https://doi.org/10.1016/S0953-7562(09)80409-7)
- Robert V, Vu D, Ben Hadj Amor B et al. (2013) MycoBank gearing up for new horizons. *IMA Fungus* 4: 371–379. <https://doi.org/10.5598/imafungus.2013.04.02.16>
- Roberts P (2000) Corticioid fungi from Korup National Park, Cameroon. *Kew Bulletin* 55: 803–842. <https://doi.org/10.2307/4113628>
- Rodríguez F, Oliver JF, Martín A, Medina JR (1990) The general stochastic model of nucleotide substitution. *Journal of Theoretical Biology* 142: 485–501. [https://doi.org/10.1016/S0022-5193\(05\)80104-3](https://doi.org/10.1016/S0022-5193(05)80104-3)
- Ronquist F, Teslenko M, Van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61: 539–542. <https://doi.org/10.1093/sysbio/sys029>
- Ryvarden L, Hjortstam K, Iturriaga T (2005) Studies in corticioid fungi from Venezuela II (Basidiomycotina, Aphyllophorales). *Synopsis Fungorum* 20: 42–78.
- Shear CL (1925) The life history of the Texas root rot fungus *Ozonium omnivorum* Shear. *Journal of Agricultural Research* 30: 475–477.
- Sjökvist E, Larsson E, Eberhardt U, Ryvarden L, Larsson KH (2012) Stipitate stereoid basidiocarps have evolved multiple times. *Mycologia* 104: 1046–1055. <https://doi.org/10.3852/11-174>
- Swofford DL (2003) PAUP\*. Phylogenetic analysis using parsimony (\*and other methods) Version 4. Sinauer Associates, Sunderland, Massachusetts.

- Telleria MT, Dueñas M, Melo I, Hallenberg N, Martín MP (2010) A re-evaluation of *Hypoch-nicium* (Polyporales) based on morphological and molecular characters. *Mycologia* 102: 1426–1436. <https://doi.org/10.3852/09-242>
- Thorn RG, Moncalvo J-M, Reddy CA, Vilgalys R (2000) Phylogenetic analyses and the distribution of nematophagy support a monophyletic Pleurotaceae within the polyphyletic pleurotoid-lentinoid fungi. *Mycologia* 92: 241–252. <https://doi.org/10.2307/3761557>
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. *Journal of Bacteriology* 172: 4238–4246. <https://doi.org/10.1128/jb.172.8.4238-4246.1990>
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gelfand DH, Sninsky JJ, White TJ (Eds) *PCR protocols: a guide to methods and applications*. Academic Press, USA, 315–322. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wilson AW, Binder M, Hibbett DS (2012) Diversity and evolution of ectomycorrhizal host associations in the Sclerodermatineae (Boletales, Basidiomycota). *New Phytologist* 194: 1079–1095. <https://doi.org/10.1111/j.1469-8137.2012.04109.x>
- Wu SH, Nilsson HR, Chen CT, Yu SY, Hallenberg N (2010) The white-rotting genus *Phanerochaete* is polyphyletic and distributed throughout the phleboid clade of Polyporales (Basidiomycota). *Fungal Diversity* 42: 107–118. <https://doi.org/10.1007/s13225-010-0031-7>.